

Symmetric Pascal matrices modulo p

Roland Bacher* and Robin Chapman

30 January 2003

1 Introduction

This paper presents results and conjectures concerning symmetric matrices associated to Pascal's triangle. We first give a formula for the determinant over $\mathbf{Z}$ of the reduction modulo 2 with values in $\{0, 1\}$ for such a matrix. We then study the reduction modulo a prime p of the characteristic polynomials of these matrices. Our main results imply a formula for the prime $p = 2$ and a conjectural formula for $p = 3$.

Consider the symmetric matrix $P(n)$ with coefficients

$$p_{i,j} = \binom{i+j}{i}, \quad 0 \leq i, j < n.$$

We call $P(n)$ the *symmetric Pascal matrix* of order n . The entries of $P(n)$ satisfy the recurrence

$$p_{i,j} = p_{i-1,j} + p_{i,j-1}.$$

In [2] the first author studied the determinant of the general matrix with entries satisfying this recurrence.

An easy computation yields $P(\infty) = T T^t$ where T is the infinite unipotent lower triangular matrix

$$T = \begin{pmatrix} 1 & & & & \\ 1 & 1 & & & \\ 1 & 2 & 1 & & \\ 1 & 3 & 3 & 1 & \\ \vdots & & & & \ddots \end{pmatrix} = \exp \begin{pmatrix} 0 & & & & \\ 1 & 0 & & & \\ 0 & 2 & 0 & & \\ & 0 & 3 & 0 & \\ & & & & \ddots \end{pmatrix}$$

with coefficients $t_{i,j} = \binom{i}{j}$. This shows that $\det(P(n)) = 1$ and that $P(n)$ is positive definite for all $n \in \mathbf{N}$. Hence all zeroes of the characteristic polynomial $\chi_n(t) = \det(tI(n) - P(n))$ (where $I(n)$ denotes the identity

*Support from the Swiss National Science Foundation is gratefully acknowledged.

matrix of size n) of $P(n)$ are positive reals. The inverse $P(n)^{-1}$ of $P(n)$ is given by

$$P(n)^{-1} = \left(T(n)^t\right)^{-1} T(n)^{-1}$$

and $T(n)^{-1}$ has coefficients $(-1)^{i+j} \binom{i}{j}$, $0 \leq i, j < n$. Hence $T(n)$ and $T(n)^{-1}$ are conjugate, and thus also $P(n)$ and $P(n)^{-1}$ are conjugate. The characteristic polynomial $\chi_n(t)$ therefore satisfies $\chi_n(t) = (-t)^n \chi(1/t)$ and 1 is always an eigenvalue of $P(2n+1)$, cf. [4]. The polynomials $\chi_n(t)$, especially their behaviour modulo primes, will be our main object of study. For convenience, we write I for $I(n)$ whenever the size of the identity matrix is unambiguous.

Define $\bar{P}(n)_2$ as the reduction modulo 2 of $P(n)$ with values in $\{0, 1\}$ by setting

$$\bar{p}_{i,j} = \left(\binom{i+j}{i} \pmod{2} \right) \in \{0, 1\}.$$

The Thue-Morse sequence $s_n = \sum \nu_i \pmod{2}$ counts the parity of all non-zero digits of a binary integer $n = \sum \nu_i 2^i$. It can also be defined recursively by $s_0 = 0$, $s_{2k} = s_k$ and $s_{2k+1} = 1 - s_k$ (cf. for instance [1]).

Theorem 1.1 *The determinant (over $\mathbf{Z}$) of $\bar{P}(n)_2$ is given by*

$$\det(\bar{P}(n)_2) = \prod_{k=0}^{n-1} (-1)^{s_k}.$$

A similar result holds for the reduction modulo 3 of $P(n)$ with values in $\{-1, 0, 1\}$.

In the sequel, we will be interested in the characteristic polynomial $\det(tI - P(n)) \pmod{p}$ for p a prime number. The next result yields a formula for $n = p^l$ and is of crucial importance in the sequel.

Proposition 1.2 *Given a power $q = p^l$ of a prime p , the matrix $P(q)$ has order 3 over $\mathbf{F}_p$. Its characteristic polynomial $\chi_q(t) = \det(tI(q) - P(q))$ satisfies*

$$\chi_q(t) \equiv (t^2 + t + 1)^{\frac{q-\epsilon(q)}{3}} (t - 1)^{\frac{q+2\epsilon(q)}{3}} \pmod{p}$$

where $\epsilon(q) \in \{-1, 0, 1\}$ satisfies $\epsilon(q) \equiv q \pmod{3}$.

In particular, $P(q)$ can be diagonalized over $\mathbf{F}_{p^2}$ except when $p = 3$. For instance, $P(3)$ has a unique Jordan block over $\mathbf{F}_3$.

This proposition (except for the diagonalization part) admits the following generalization:

Theorem 1.3 *When $q = p^l$ is a power of a prime p and $0 \leq k \leq q/2$ then*

$$\chi_{q-k}(t) \equiv (t^2 + t + 1)^{(q-\epsilon(q))/3-k} (t - 1)^{(q+2\epsilon(q))/3-k} \det(t^2 I + P(k)) \pmod{p}$$

where $\epsilon(q) \in \{-1, 0, 1\}$ satisfies $\epsilon(q) \equiv q \pmod{3}$.

Theorem 1.3 completely determines the reduction modulo 2 of $\chi_n(t)$ as follows: Define a sequence $\gamma(0) = 0, \gamma(1), \dots$ recursively by

$$\gamma(2^l - k) = \frac{2^l + 2(-1)^l}{3} - k + 2\gamma(k), \quad 0 \leq k \leq 2^{l-1}.$$

Theorem 1.4 *For all $n \in \mathbf{N}$*

$$\chi_n(t) \equiv (t+1)^{\gamma(n)}(t^2+t+1)^{\gamma_2(n)} \pmod{2}$$

where $\gamma_2(n) = \frac{1}{2}(n - \gamma(n))$.

It follows immediately that the matrix $I - P(n)^3$ is nilpotent over $\mathbf{F}_2$ for all $n \in \mathbf{N}$.

The first terms $\gamma(1), \dots, \gamma(32)$ and $\gamma_2(1), \dots, \gamma_2(32)$ are given by

n	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$\gamma(n)$	1	0	3	2	5	0	3	2	5	0	11	6	9	4	7	6
$\gamma_2(n)$	0	1	0	1	0	3	2	3	2	5	0	3	2	5	4	5
n	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32
$\gamma(n)$	9	4	15	10	21	0	11	6	9	4	15	10	13	8	11	10
$\gamma_2(n)$	4	7	2	5	0	11	6	9	8	11	6	9	8	11	10	11

The sequence $\gamma(0), \gamma(1), \dots$ has many interesting arithmetic features. In order to describe them, let us introduce the number $b(n)$ of “blocks” of adjacent ones in the binary representation of a positive integer n . For instance $667 = (1010011011)_2$ and so $b(667) = 4$. Notice that $b(2n) = b(n)$ and $b(2n+1) = b(n) + 1 - (n \pmod{2})$ (with $n \pmod{2} \in \{0, 1\}$). This, together with $b(0) = 0$, defines the sequence $b(n)$ recursively.

Theorem 1.5 (i) *We have*

$$\gamma(2^l + k) = \frac{2^l + 2(-1)^l}{3} - k + 4\gamma(k)$$

for all $0 \leq k \leq 2^{l-1}$.

(ii) *We have for all $n \in \mathbf{N}$ and $2^{l-2} \leq k \leq 2^{l-1}$*

$$\gamma(2^l - k) = \gamma(k) + 2\gamma(2^{l-1} - k).$$

(iii) *We have*

$$\gamma(2^l + k) = 1 + \gamma(2^l + k - 1) + 2\gamma(2^l - k) - 2\gamma(2^l + 1 - k)$$

for $1 \leq k \leq 2^l$.

(iv) *We have*

$$\begin{aligned} \gamma(2n) &= n - \gamma(n), \\ \gamma(2n-1) &= \gamma(2n) + (4^{b(2n-1)} - 1)/3 = n - \gamma(n) + (4^{b(2n-1)} - 1)/3, \\ \gamma(2n+1) &= \gamma(2n) + (2^{1+2b(n)} + 1)/3 = n - \gamma(n) + (2^{1+2b(n)} + 1)/3. \end{aligned}$$

Part (iv) of this Theorem gives an alternative recursive definition of the sequence $(\gamma(n))$.

Theorem 1.3 seems to have many generalizations. A first one is given by the following:

Conjecture 1.6 *For each integer $k \geq 0$ there exists a monic polynomial $c_k(t) \in \mathbf{Z}[t]$ of degree $4k$ such that $c_k(t) = t^{4k}c_k(t^{-1})$ with the following property: if q is a power of a prime p , and $0 \leq k \leq q/2$ then*

$$\chi_{q+k}(t) \equiv (t^2 + t + 1)^{(q-\epsilon(q))/3-k} (t-1)^{(q+2\epsilon(q))/3-k} c_k(t) \pmod{p}$$

where $\epsilon(q) \in \{-1, 0, 1\}$ satisfies $\epsilon(q) \equiv q \pmod{3}$.

The first few of these conjectural polynomials $c_k(t)$ are

$$\begin{aligned} c_0(t) &= 1, \\ c_1(t) &= t^4 - 2t^3 - 2t + 1, \\ c_2(t) &= t^8 - 6t^7 + 4t^6 - 4t^5 + 15t^4 - 4t^3 + 4t^2 - 6t + 1, \\ c_3(t) &= (t^4 - 2t^3 - 2t + 1)(t^8 - 16t^7 + 4t^6 - 4t^5 + 40t^4 - 4t^3 + 4t^2 - 16t + 1), \\ c_4(t) &= t^{16} - 58t^{15} + 288t^{14} - 240t^{13} + 393t^{12} - 1440t^{11} + 836t^{10} - 902t^9 \\ &\quad + 2376t^8 - 902t^7 + \cdots - 58t + 1, \\ c_5(t) &= c_1(t)(t^{16} - 196t^{15} + 2112t^{14} - 792t^{13} + 1290t^{12} - 10560t^{11} \\ &\quad + 2768t^{10} - 2972t^9 + 17424t^8 - 2972t^7 + \cdots - 196t + 1). \end{aligned}$$

For $p = 2$, it follows from Theorem 1.4 and assertion (ii) in Theorem 1.5 that if $c_k(t)$ exists then

$$c_k(t) \equiv (\det(tI + P(k)))^4 \pmod{2}.$$

Computations suggest:

Conjecture 1.7 *We have*

$$c_k(t) \equiv (t+1)^{3k} \det(tI + P(k)) \pmod{3}.$$

This conjecture, together with Theorem 1.3 yields conjectural recursive formulas for $p_n(t) = \det(tI(n) - P(n)) \pmod{3}$ as follows: Set $p_0(t) = 1 \pmod{3}$, $p_1(t) = 1 - t \pmod{3}$. For $n = 3^l \pm k > 1$ with $0 \leq k < \frac{3^l}{2}$ the characteristic polynomial $\chi_n(t) \pmod{3}$ is then conjecturally given by

$$\begin{aligned} (t-1)^{3^l-3k} \det(t^2I + P(k)) &\quad \text{if } n = 3^l - k, \\ (t-1)^{3^l-3k} (t+1)^{3k} \det(tI + P(k)) &\quad \text{if } n = 3^l + k. \end{aligned}$$

In particular, all roots of $\chi_n(t)$ modulo 3 should be of multiplicative order a power of 2 in the algebraic closure of $\mathbf{F}_3$.

We conclude finally by mentioning a last conjectural observation:

Conjecture 1.8 *Given a prime-power $q = p^l \equiv 2 \pmod{3}$, we have*

$$\chi_{(q+1)/3}(t) \equiv (t+1)^{(q+1)/3} \pmod{p}$$

and

$$\chi_{(2q-1)/3}(t) \equiv (t+1)^{(q+1)/3} (t-1)^{(q-2)/3} \pmod{p}.$$

Remark 1.9 (i) *The matrix $C = P(\frac{q+1}{3}) + I(\frac{q+1}{3})$ for $q = p^l \equiv 2 \pmod{3}$ a prime-power, appears to have a unique Jordan block of maximal length over $\mathbf{F}_p$. If so, the rows of $C^{(q+1)/6}$ generate a self-dual code over $\mathbf{F}_p$.*

(ii) *Given a prime power $q = p^l \equiv 2 \pmod{3}$ as above we set $n = \frac{2q+2}{3}$ and $k = \frac{2q-1}{3}$. We conjecture that the characteristic polynomial of the matrix $\tilde{P}_k(n)$ with coefficients*

$$\tilde{p}_{i,j} = \binom{i+j+2k}{i+k}, \quad 0 \leq i, j < n$$

satisfies $\det(tI - \tilde{P}_k(n)) \equiv (1+t)^n \pmod{p}$.

Remark 1.10 *In [3, Theorems 32 and 35] Krattenthaler gives evaluations of determinants related to ours, namely of $\det(\omega I + Q(n))$ where ω is a sixth root of unity, and $Q(n)$ has entries $\binom{2\mu+i+j}{j} (0 \leq i, j < n)$.*

The sequel of this paper is organized as follows:

Section 2 is devoted to autosimilar matrices. Such matrices generalize the matrix $\overline{P}(\infty)_2$ and their properties imply easily Theorem 1.1.

Section 3 contains proofs of Proposition 1.2 and Theorem 1.3.

Section 4 contains proofs of Theorems 1.4 and 1.5.

2 Autosimilar matrices

Let $b \geq 1$ be a natural integer. An infinite matrix M with coefficients $m_{i,j}$ ($i, j \geq 0$) is *b-autosimilar* if $m_{0,0} = 1$ and if

$$m_{s,t} = \prod_i m_{\sigma_i, \tau_i}$$

where the indices $s = \sum \sigma_i b^i$, $t = \sum \tau_i b^i$ are written in base b , that is, $\sigma_i, \tau_i \in \{0, \dots, b-1\}$ for all $i = 0, 1, 2, \dots$

We denote by $M(n)$ the finite sub-matrix of M with coefficients $m_{i,j}$, $0 \leq i, j < n$. A b -autosimilar matrix M is *non-degenerate* if the determinants

$$\det(M(n))$$

are invertible for $n = 2, \dots, b$.

Theorem 2.1 *Let $b \geq 2$ be an integer and let M be a b -autosimilar matrix which is non-degenerate. One has then a factorization*

$$M = LDU$$

where L, D, U are b -autosimilar and where L is unipotent lower-triangular, D is diagonal and U is unipotent upper-triangular.

Corollary 2.2 *Given a non-degenerate b -autosimilar matrix M one has*

$$\det(M(n)) = \prod_{i=0}^{n-1} d_{\nu_i}$$

for all $n = \sum \nu_i b^i$ with $d_0 = 1$ and

$$d_k = \det(M(k+1)) / \det(M(k))$$

for $k = 1, \dots, b-1$.

Remark 2.3 *In general, one can compute determinants of arbitrary b -autosimilar matrices over a field K by applying Corollary 2.2 to the b -autosimilar matrix obtained from a generic perturbation of the form*

$$M_t(b) = (1-t)M(b) + tP(b)$$

(where $P(b)$ is a suitable matrix) and working over the rational function field $K(t)$.

Proof of Theorem 2.1. The genericity of M implies that

$$M(b) = L(b)D(b)U(b)$$

where $L(b)$ and $U(b)$ are unipotent upper and lower triangular matrices and the diagonal matrix $D(b)$ has entries $d_{0,0} = 1$ and $d_{k,k} = \det(M(k+1)) / \det(M(k))$ for $k = 1, \dots, b-1$. Extending $L(b)$, $D(b)$ and $U(b)$ in the unique possible way to infinite b -autosimilar matrices L , D and U we have

$$\begin{aligned} (LDU)_{s,t} &= \sum_k L_{s,k} D_{k,k} U_{k,t} \\ &= \sum_{k=\sum \kappa_i b^i} \prod_i L_{\sigma_i, \kappa_i} D_{\kappa_i, \kappa_i} U_{\kappa_i, \tau_i} \\ &= \prod_i \sum_{\kappa_i=0}^{b-1} L_{\sigma_i, \kappa_i} D_{\kappa_i, \kappa_i} U_{\kappa_i, \tau_i} \\ &= \prod_i M_{\sigma_i, \tau_i} = M_{s,t} \end{aligned}$$

for all $s = \sum \sigma_i b^i, t = \sum \tau_i b^i \in \mathbb{N}$. □

The identity

$$\det(M(n)) = \det(D(n))$$

implies immediately Corollary 2.2.

2.1 Binomial coefficients modulo a prime p

Let p be a prime number. We have then

$$(1+x)^n = \prod (1+x)^{\nu_i p^i} \equiv (1+x^{p^i})^{\nu_i} \pmod{p}$$

(using properties of the Frobenius automorphism in characteristic p). This implies immediately the equality

$$\binom{n}{k} = \prod_i \binom{\nu_i}{\kappa_i}$$

allowing (for small primes) an efficient computation of binomial coefficients $\pmod{p}$.

This equality shows that the reductions modulo 2 or 3 of the symmetric Pascal triangle P with coefficients

$$\bar{p}_{i,j} = \left(\binom{i+j}{i} \pmod{2} \right) \in \{0,1\}$$

respectively

$$\bar{p}_{i,j} = \left(\binom{i+j}{i} \pmod{3} \right) \in \{-1,0,1\}$$

are 2– (respectively 3–) autosimilar matrices.

For $p = 2$ we have

$$\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

which yields $d_0 = 1, d_1 = -1$ and Corollary 2.2 implies now Theorem 1.1.

Remark 2.4 *One can show that the inverse of the integral matrix $\bar{P}(n)_2$ considered in Theorem 1.1 has all its coefficients in $\{-1,0,1\}$ for all n .*

For $p = 3$ we have

$$\begin{pmatrix} 1 & 1 & 1 \\ 1 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & \frac{1}{2} & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & \frac{1}{2} \\ 0 & 0 & 1 \end{pmatrix}$$

This shows that $\det(\bar{P}(n)_3)$ (over $\mathbf{Z}$) equals $(-2)^{a-b}$ where a and b are the number of digits 1 and 2 needed in order to write all natural integers $< n$ in base 3.

3 Proofs of Proposition 1.2 and Theorem 1.3

Proof of Proposition 1.2 Let R be a commutative ring, and let

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}(2, R).$$

Then A determines a (graded R -algebra) automorphism ϕ_A of $R[X, Y]$ via $\phi_A(X) = aX + bY$ and $\phi_A(Y) = cX + dY$, or alternatively

$$\begin{pmatrix} \phi_A(X) \\ \phi_A(Y) \end{pmatrix} = A \begin{pmatrix} X \\ Y \end{pmatrix}.$$

It is easy to see that $\phi_A \circ \phi_B = \phi_{BA}$. Each ϕ_A restricts to an R -module automorphism of the homogeneous polynomials $R[X, Y]_{n-1}$ of degree $n-1$. Let $A^{(n)}$ denote the matrix of this endomorphism with respect to the basis $X^{n-1}, X^{n-2}Y, X^{n-3}Y^2, \dots, Y^{n-1}$, that is

$$\begin{pmatrix} \phi_A(X^{n-1}) \\ \phi_A(X^{n-2}Y) \\ \phi_A(X^{n-3}Y^2) \\ \vdots \\ \phi_A(Y^{n-1}) \end{pmatrix} = A^{(n)} \begin{pmatrix} X^{n-1} \\ X^{n-2}Y \\ X^{n-3}Y^2 \\ \vdots \\ Y^{n-1} \end{pmatrix}.$$

Then $A^{(n)} \in \mathrm{GL}(n, R)$ and $(AB)^{(n)} = A^{(n)}B^{(n)}$. (Another way of expressing this is to say that $A^{(n)}$ is the $(n-1)$ -th symmetric power of A .)

Let us specialize to the case $R = \mathbf{F}_p = \mathbf{Z}/p\mathbf{Z}$ and $n = p^l$. In this case $A^{(n)} = I$ if and only if A is a scalar matrix. The matrix

$$A = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}$$

yields $A^{(n)} \equiv P(p^l) \pmod{p}$. Since $A^3 = -I$, the matrix $A^{(n)}$ has order 3.

Let us now compute the multiplicities of the three eigenvalues of $P = P(p) \pmod{p}$ over $\mathbf{F}_p$ (the formula for $P(p^l)$ is then a straightforward consequence of the fact the $P(p^l)$ is the l -fold Kronecker product of $P(p)$ with itself).

The easy identity $\binom{2k}{k} = \binom{(p-1)/2}{k}(-4)^k \pmod{p}$ for p an odd prime and $0 \leq k \leq (p-1)/2$ shows

$$\sum_{k=0}^{(p-1)/2} \binom{2k}{k} \left(\frac{-x}{4}\right)^k \equiv (1+x)^{(p-1)/2} \pmod{p}$$

and yields $\mathrm{tr}(P) \equiv (-3)^{(p-1)/2} \equiv \epsilon(p) \pmod{p}$ (where $\epsilon(p) \in \{-1, 0, 1\}$ satisfies $\epsilon(p) \equiv p \pmod{3}$) by quadratic reciprocity.

Since the characteristic polynomial for P has antisymmetric coefficients ($\alpha_k = -\alpha_{p-k}$) the two eigenvalues $\neq 1$ of P have equal multiplicity r . Lifting into positive integers $\leq \frac{p-1}{2}$ the solution of the linear system $-r + (p-2r) \equiv \text{tr}(P) \pmod{p}$ yields now the result.

The case $p = 2$ is easily solved by direct inspection. $\square$

Remark 3.1 Recall that we have (with the notations of the above proof) $P = P(n) = A^{(n)} \pmod{p}$ for $n = p^l$ and introduce $L = L(n) = B^{(n)} \pmod{p}$ and $\tilde{L} = \tilde{L}(n) = C^{(n)} \pmod{p}$ where

$$A = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}, B = \begin{pmatrix} 1 & 0 \\ -1 & -1 \end{pmatrix}, C = \begin{pmatrix} 1 & 0 \\ 1 & -1 \end{pmatrix}.$$

It is straightforward to check that L and $\tilde{L}$ have coefficients

$$l_{i,j} = (-1)^i \binom{i}{j} \pmod{p} \quad \text{and} \quad \tilde{l}_{i,j} = (-1)^j \binom{i}{j} \pmod{p}$$

for $0 \leq i, j < n$.

Then $A^3 = -I$, but $(-I)^{(n)}$ is the identity. Hence $P^3 = I$. Also $C^2 = I$ and $CAC = A^{-1}$. It follows that A and C generate a dihedral group of order 12, containing $-I$. Hence $A^{(n)} = P$ and $C^{(n)} = \tilde{L}$ generate a dihedral group of order 6.

The group G_p generated by P and L depends on the prime p (but not on the power l of $n = p^l$). It is isomorphic to a subgroup of $\text{PGL}_2(\mathbf{F}_p)$. For all but finitely many primes p , G_p is isomorphic to $\text{PSL}_2(\mathbf{F}_p)$ or $\text{PGL}_2(\mathbf{F}_p)$ according to whether -1 is or is not a square in $\mathbf{F}_p$. The exceptional primes are 5, 7 and 29 where G_p has order 24, 42 and 120 respectively.

Proof of Theorem 1.3 Using Proposition 1.2, we can rewrite the equation to be proved as

$$(t^3 - 1)^k \det(tI - P(q - k)) \equiv \det(tI - P(q)) \det(t^2I + P(k)) \pmod{p}.$$

Here, and in the sequel, we write I for $I(n)$ whenever this notation is unambiguous; also we denote the zero matrix of any size by O .

We now work over the field $\mathbf{F}_p$. Unless otherwise stated vectors will be row vectors.

It is convenient to define a category $\mathcal{E} = \mathcal{E}_{\mathbf{F}_p}$ as follows. Its objects will be pairs (V, α) where V is a finite-dimensional vector space over $\mathbf{F}_p$ and α is a vector space endomorphism of V . A morphism $\phi : (V, \alpha) \rightarrow (W, \beta)$ in $\mathcal{E}$ will be a linear map $\phi : V \rightarrow W$ with $\phi \circ \alpha = \beta \circ \phi$. (In fact $\mathcal{E}$ is equivalent to the category of finitely generated torsion modules over the polynomial ring $\mathbf{F}_p[X]$.) If (V, α) is an object of $\mathcal{E}$ we define $\chi(V, \alpha, t)$ as the characteristic polynomial of α acting on V , that is, $\chi(V, \alpha, t) = \det(tI - A)$

where A is a matrix representing α with respect to some basis of V . An r by r matrix A defines an object $((\mathbf{F}_p)^r, \alpha)$, denoted by $((\mathbf{F}_p)^r, A)$, where α is the endomorphism defined by A .

It is easy to see that $\mathcal{E}$ is an abelian category, and that if

$$0 \rightarrow (V, \alpha) \rightarrow (X, \gamma) \rightarrow (W, \beta) \rightarrow 0$$

is a short exact sequence, then $\chi(X, \gamma, t) = \chi(V, \alpha, t)\chi(W, \beta, t)$. This is because there is a basis for X with respect to which the matrix of γ (acting on row vectors from the the right) is

$$\begin{pmatrix} A & O \\ C & B \end{pmatrix}$$

where A and B are matrices representing α and β respectively.

Set $k' = q - k$. We can partition the Pascal matrices $P(k')$ and $P(q)$ as follows:

$$P(k') = \begin{pmatrix} A & B \\ B^t & C \end{pmatrix} \quad \text{and} \quad P(q) = \begin{pmatrix} A & B & D \\ B^t & C & O \\ D^t & O & O \end{pmatrix}$$

where $A = P(k)$.

Let $\overline{A}$ denote the matrix obtained by rotating A through 180° . Then $P(q)^2 = \overline{P(q)}$ and $P(q)^3 = I$. Hence

$$P(q)^2 = \begin{pmatrix} O & O & \overline{D^t} \\ O & \overline{C} & \overline{B^t} \\ \overline{D} & \overline{B} & \overline{A} \end{pmatrix}.$$

Thus

$$A^2 + BB^t + DD^t = O$$

and so

$$P(k')^2 = \begin{pmatrix} -DD^t & O \\ O & \overline{C} \end{pmatrix}.$$

From $P(q)^2 = \overline{P(q)}$ it follows that $AD = \overline{D^t}$ and from $\overline{P(q)}P(q) = I$ it follows that $\overline{D^t}D^t = I$. Hence $ADD^t = I$ and so

$$P(k')^2 = \begin{pmatrix} -A^{-1} & O \\ O & \overline{C} \end{pmatrix}.$$

Let $V = (\mathbf{F}_p)^q$ and $X = (\mathbf{F}_p)^{3k}$. Let

$$Q_1 = \begin{pmatrix} O & I(k) & O \\ O & O & I(k) \\ I(k) & O & O \end{pmatrix}.$$

Let $\phi : X \rightarrow V$ be the map defined by the matrix

$$\begin{pmatrix} I & O & O \\ A & B & D \\ O & O & \overline{D^t} \end{pmatrix}.$$

Then

$$Q_1 \begin{pmatrix} I & O & O \\ A & B & D \\ O & O & \overline{D^t} \end{pmatrix} = \begin{pmatrix} A & B & D \\ O & O & \overline{D^t} \\ I & O & O \end{pmatrix}$$

and

$$\begin{pmatrix} I & O & O \\ A & B & D \\ O & O & \overline{D^t} \end{pmatrix} P(q) = \begin{pmatrix} I & O & O \\ A & B & D \\ O & O & \overline{D^t} \end{pmatrix} \begin{pmatrix} A & B & D \\ B^t & C & O \\ D^t & O & O \end{pmatrix} = \begin{pmatrix} A & B & D \\ O & O & \overline{D^t} \\ I & O & O \end{pmatrix}$$

where we have used the formulas $P(q)^2 = \overline{P(q)}$ and $\overline{P(q)}P(q) = I$. Hence ϕ is a morphism from $((\mathbf{F}_p)^{3k}, Q_1)$ to $((\mathbf{F}_p)^q, P(q))$ in $\mathcal{E}$.

Let $W = (\mathbf{F}_p)^{k'}$ and $Y = (\mathbf{F}_p)^{2k}$. Let

$$Q_2 = \begin{pmatrix} O & I(k) \\ -A^{-1} & O \end{pmatrix}.$$

Let $\psi : Y \rightarrow W$ be the map defined by the matrix

$$\begin{pmatrix} I & O \\ A & B \end{pmatrix}.$$

Then

$$Q_2 \begin{pmatrix} I & O \\ A & B \end{pmatrix} = \begin{pmatrix} A & B \\ -A^{-1} & O \end{pmatrix}$$

and

$$\begin{pmatrix} I & O \\ A & B \end{pmatrix} P(k') = \begin{pmatrix} I & O \\ A & B \end{pmatrix} \begin{pmatrix} A & B \\ B^t & C \end{pmatrix} = \begin{pmatrix} A & B \\ -A^{-1} & O \end{pmatrix}$$

where we have used the formula

$$P(k')^2 = \begin{pmatrix} -A^{-1} & O \\ O & \overline{C} \end{pmatrix}.$$

Hence ψ is a morphism from $((\mathbf{F}_p)^{2k}, Q_2)$ to $((\mathbf{F}_p)^{k'}, P(k'))$ in $\mathcal{E}$.

We need to divide into the cases $k \leq q/3$ and $k \geq q/3$. In the former cases ϕ and ψ are injective and in the latter case they are surjective. In the former case we consider their cokernels, in the latter case their kernels.

The matrix B has size k by $q-2k$. If B has rank k (which is only possible if $k \leq q/3$) then ϕ and ψ are injective. If B has rank $q-2k$ (which is only possible if $k \geq q/3$) then ϕ and ψ are surjective.

The matrix B contains a submatrix

$$\left(\binom{i+j+k}{i} \right)_{i,j=0}^{r-1}$$

where $r = \min(k, q-2k)$. This submatrix has determinant 1 (consider it as a matrix over $\mathbf{Z}$ and reduce it to a Vandermonde matrix or see for instance [2]). Thus B has rank r and indeed ϕ and ψ are injective for $k \leq q/3$ and surjective for $k \geq q/3$.

Consider first the case where $k \leq q/3$. Let (X_1, θ_1) and (X_2, θ_2) denote the cokernels of $\phi : ((\mathbf{F}_p)^{3k}, Q_1) \rightarrow ((\mathbf{F}_p)^q, P(q))$ and $\psi : ((\mathbf{F}_p)^{2k}, Q_2) \rightarrow ((\mathbf{F}_p)^{k'}, P(k'))$ in $\mathcal{E}$. Then

$$\chi((\mathbf{F}_p)^q, P(q), t) = \chi((\mathbf{F}_p)^{3k}, Q_1, t) \chi(X_1, \theta_1, t)$$

and

$$\chi((\mathbf{F}_p)^{k'}, P(k'), t) = \chi((\mathbf{F}_p)^{2k}, Q_2, t) \chi(X_2, \theta_2, t).$$

It is apparent that

$$\chi((\mathbf{F}_p)^{3k}, Q_1, t) = (t^3 - 1)^k$$

and

$$\chi((\mathbf{F}_p)^{2k}, Q_2, t) = \det(t^2 I + A^{-1}) = \det(t^2 I + A)$$

as A and A^{-1} are similar. Hence

$$\det(tI - P(q)) = (t^3 - 1)^k \chi(X_1, \theta_1, t)$$

and

$$\det(tI - P(k')) = \det(t^2 I + A) \chi(X_2, \theta_2, t).$$

It suffices to prove that (X_1, θ_1) and (X_2, θ_2) are isomorphic in $\mathcal{E}$.

As $\overline{D}^t$ is nonsingular, it is apparent that X_1 is isomorphic to $(\mathbf{F}_p)^{q-2k}/Y$ where Y is the row space of B and that the action of θ_1 is induced by that of the matrix C on $(\mathbf{F}_p)^{q-2k}$. It is even more apparent that X_2 is isomorphic to $(\mathbf{F}_p)^{q-2k}/Y$ and that the action of θ_2 is induced by C . Hence (X_1, θ_1) and (X_2, θ_2) are isomorphic in $\mathcal{E}$. This completes the argument in the case $k \leq q/3$.

Now suppose that $k \geq q/3$. Let (K_1, θ_1) and (K_2, θ_2) denote the kernels of $\phi : ((\mathbf{F}_p)^{3k}, Q_1) \rightarrow ((\mathbf{F}_p)^q, P(q))$ and $\psi : ((\mathbf{F}_p)^{2k}, Q_2) \rightarrow ((\mathbf{F}_p)^{k'}, P(k'))$ in $\mathcal{E}$. Then

$$\chi((\mathbf{F}_p)^q, P(q), t) \chi(K_1, \theta_1, t) = \chi((\mathbf{F}_p)^{3k}, Q_1, t)$$

and

$$\chi((\mathbf{F}_p)^{k'}, P(k'), t) \chi(K_2, \theta_2, t) = \chi((\mathbf{F}_p)^{2k}, Q_2, t).$$

Hence

$$\frac{(t^3 - 1)^k}{\det(tI - P(q))} = \chi(K_1, \theta_1, t)$$

and

$$\frac{\det(t^2 I + A)}{\det(tI - P(k'))} = \chi(K_2, \theta_2, t).$$

It suffices to prove that (K_1, θ_1) and (K_2, θ_2) are isomorphic in $\mathcal{E}$.

As $\overline{D^t}$ is nonsingular and has inverse D^t , it is apparent that

$$K_1 = \{(-uA, u, -uDD^t) = (-uA, u, -uA^{-1}) : u \in (\mathbf{F}_p)^k, uB = 0\}$$

and we have

$$(-uA, u, -uA^{-1})Q_1 = (-uA^{-1}, -uA, u) .$$

Also

$$K_2 = \{(-uA, u) : u \in (\mathbf{F}_p)^k, uB = 0\}$$

and

$$(-uA, u)Q_2 = (-uA^{-1}, -uA) .$$

Hence the linear map

$$(-uA, u, -uA^{-1}) \mapsto (-uA, u)$$

induces an isomorphism between (K_1, θ_1) and (K_2, θ_2) . $\square$

4 Proofs for the prime $p = 2$

Proof of Theorem 1.4. Set $n = 2^l - k$ and $q = 2^l$ where $1 \leq k \leq 2^{l-1}$.

Theorem 1.3 yields then over $\mathbf{F}_2$

$$\chi_n(t) = \chi_{q-k}(t) = (t^2 + t + 1)^{(q-\epsilon(q))/3-k} (t+1)^{(q+2\epsilon(q))/3-k} \det(tI + P(k))^2$$

since $x \mapsto x^2$ is an automorphism in characteristic 2.

By induction on l , the only possible irreducible factors of $\det(tI(n) - P(n)) \pmod{2}$ are $(1+t)$ and $(1+t+t^2)$. The multiplicity $\mu(n) = \mu(2^l - k)$ of the factor $(1+t)$ in this polynomial is hence recursively defined by

$$\mu(n) = \frac{2^l + 2(-1)^l}{3} - k + 2\mu(k)$$

and coincides hence with the sequence γ of Theorem 1.4. The remaining factor of $\det(tI(n) - P(n)) \pmod{2}$ is hence given by $(1+t+t^2)^{\gamma_2(n)}$ where $\gamma_2(n) = \frac{1}{2}(n - \gamma(n))$ and this proves the result. $\square$

Proof of Theorem 1.5. We have for $0 \leq k \leq 2^{l-1}$

$$\begin{aligned}
\gamma(2^l + k) &= \gamma(2^{l+1} - (2^l - k)) \\
&= \frac{2^{l+1} - 2(-1)^l}{3} - 2^l + k + 2\gamma(2^l - k) \\
&= \frac{2^{l+1} - 2(-1)^l}{3} - 2^l + k + 2 \frac{2^l + 2(-1)^l}{3} - 2k + 4\gamma(k)
\end{aligned}$$

which is assertion (i).

We have for all $2^{l-2} \leq k \leq 2^{l-1}$

$$\begin{aligned}
\gamma(2^l - k) &= \frac{2^l + 2(-1)^l}{3} - k + \gamma(k) + \gamma(2^{l-1} - (2^{l-1} - k)) \\
&= \frac{2^l + 2(-1)^l}{3} - k + \gamma(k) + \frac{2^{l-1} - 2(-1)^l}{3} - 2^{l-1} + k + 2\gamma(2^{l-1} - k) \\
&= \gamma(k) + 2\gamma(2^{l-1} - k)
\end{aligned}$$

which proves assertion (ii).

Similarly, we have for $1 \leq k \leq 2^l$

$$\begin{aligned}
\gamma(2^l + k) - \gamma(2^l + k - 1) &= \gamma(2^{l+1} - (2^l - k)) - \gamma(2^{l+1} - (2^l - k + 1)) \\
&= 1 + 2\gamma(2^l - k) - 2\gamma(2^l - k + 1)
\end{aligned}$$

which proves assertion (iii).

Writing $2n = 2^l - 2k$ with $1 \leq k \leq 2^{l-2}$ we have, using induction on n ,

$$\begin{aligned}
\gamma(2^l - 2k) &= \frac{2^l - (-1)^l}{3} - 2k + 2\gamma(2k) \\
&= \frac{2^l - (-1)^l}{3} - 2k + 2(k - \gamma(k)) \\
&= (2^{l-1} - k) - \left(\frac{2^{l-1} - (-1)^{l-1}}{3} - k + 2\gamma(k) \right) \\
&= (2^{l-1} - k) - \gamma(2^{l-1} - k)
\end{aligned}$$

which proves the first equality of assertion (iv) (this equality follows also from the fact that $P(2n)$ is the Kronecker product of $P(n)$ with $P(2)$ over $\mathbf{F}_2$).

The second identity of assertion (iv) amounts to the equality

$$\gamma(2n - 1) - \gamma(2n) = \frac{4^{b(2n-1)} - 1}{3} .$$

We prove first by induction on n that this identity is equivalent to the last identity.

The last identity and induction yield

$$\begin{aligned}\gamma(2n-1) - \gamma(2n) &= \gamma(2n-1) - \gamma(2n-2) + \gamma(2n-2) - \gamma(2n) \\ &= \frac{2^{1+2b(n-1)} + 1}{3} - 1 + \gamma(n) - \gamma(n-1).\end{aligned}$$

We now divide into cases according to the parity of n .

Suppose first that $n = 2m$ is even. Then inductively

$$\gamma(n) - \gamma(n-1) = \gamma(2m) - \gamma(2m-1) = -\frac{4^{b(2m-1)-1}}{3} = -\frac{4^{b(n-1)-1}}{3}$$

Hence

$$\gamma(2n-1) - \gamma(2n) = -1 + \frac{2^{1+2b(n-1)} + 1}{3} - \frac{2^{2b(n-1)} - 1}{3} = \frac{2^{2b(n-1)} - 1}{3}.$$

But

$$2^{2b(n-1)} = 4^{b(n-1)} = 4^{b(2n-1)}$$

as the binary representation of $n-1$ ends in 1 and that of $2n-1$ is obtained by appending 1.

Now suppose that $n = 2m+1$ is odd. Then

$$\gamma(n) - \gamma(n-1) = \gamma(2m+1) - \gamma(2m) = \frac{2^{1+2b(m)} + 1}{3} = \frac{2^{1+2b(2m)} + 1}{3}.$$

Hence

$$\gamma(2n-1) - \gamma(2n) = -1 + \frac{2^{1+2b(n-1)} + 1}{3} + \frac{2^{1+2b(n-1)} + 1}{3} = \frac{2^{2+2b(n-1)} - 1}{3}.$$

But

$$2^{2+2b(n-1)} = 4^{1+b(n-1)} = 4^{b(2n-1)}$$

as the binary representation of $n-1$ ends in 0 and that of $2n-1$ is obtained by appending 1.

This completes the proof of equivalence of the two last identities in assertion (iv).

We prove now the last identity by induction on n .

The last identity of assertion (iv) is equivalent to

$$\gamma(2n+1) - \gamma(2n) = \frac{2^{1+2b(n)} + 1}{3}.$$

Writing $2n+1 = 2^l + k$ with $1 \leq k < 2^l$ and applying assertion (iii) and the second identity of assertion (iv) (which holds by induction) we have

$$\begin{aligned}\gamma(2n+1) - \gamma(2n) &= 1 + 2\gamma(2^l - k) - 2\gamma(2^l + 1 - k) \\ &= 1 + 2\frac{4^{b(2^l-k)} - 1}{3} \\ &= \frac{2^{1+2b(2^l-k)} + 1}{3}\end{aligned}$$

Since $(2^l + k - 1) + (2^l - k) = 2^{l+1} - 1$ and since $2^l + k - 1$ is even and greater than $2^l - k$, they have the same number of blocks $1 \dots 1$ in their binary expansion. This shows $b(2^l - k) = b(2n) = b(n)$ and establishes the last identity of assertion (iv). $\square$

The first author wishes to thank J.-P. Allouche, F. Sigris, U. Vishne and A. Wassermann for interesting comments and remarks.

References

- [1] J.-P. Allouche, J. Shallit, *The ubiquitous Prouhet-Thue-Morse sequence*, Proceedings of SETA 98 (C. Ding, T. Hellese, H. Niederreiter, editors), Springer (1999).
- [2] R. Bacher, *Determinants of matrices related to the Pascal triangle*, J. de Th. des Nombres de Bordeaux **14** (2002), 19–41.
- [3] C. Krattenthaler, *Advanced determinant calculus*, Sémin. Lothar. Comb. 42, B42q (1999), 67 pages.
- [4] W.F. Lunnon, *The Pascal matrix*, Fib. Quart. vol. 15 (1977), 201–204.

Roland Bacher, Institut Fourier, UMR 5582, Laboratoire de Mathématiques,
BP 74, 38402 St. Martin d’Hères Cedex, France, Roland.Bacher@ujf-grenoble.fr

Robin Chapman, University of Exeter, School of Mathematical Sciences,
North Park Road, EX4 4QE Exeter, UK, rjc@maths.ex.ac.uk